

## Mechanisms of allergic diseases

# Innate lymphoid cells in allergic and nonallergic inflammation



Hideaki Morita, MD, PhD,<sup>a,b</sup> Kazuyo Moro, PhD,<sup>c,d</sup> and Shigeo Koyasu, PhD<sup>e,f</sup>

Davos, Switzerland, and Yokohama and

Tokyo, Japan

In the last decade, the full picture of the role of innate lymphoid cells (ILCs) has been gradually revealed. ILCs are classified into 3 groups based on their transcription factors and cytokine production patterns, which mirror helper T-cell subsets. Unlike T cells and B cells, ILCs do not have antigen receptors. They promptly respond to multiple tissue-derived factors, such as cytokines and alarmins, and produce multiple proinflammatory and immunoregulatory cytokines. It has been reported that ILC-derived cytokines are important for the induction and regulation of inflammation.

Accumulating evidence suggests that ILCs play substantial roles in protection against infection and the pathogenesis of inflammatory diseases, such as allergic diseases and autoimmune diseases.

Different ILC subsets localize in distinct tissue/organ niches and receive tissue-derived signals on different types of inflammation, which allows them to acquire diverse phenotypes with specialized effector capacities. In this review we highlight the roles of ILCs in a variety of organs, such as the airway, skin, and gastrointestinal tract, in the context of allergic and nonallergic inflammation. (*J Allergy Clin Immunol* 2016;138:1253-64.)

**Key words:** *Innate lymphoid cell, asthma, viral infection, obesity, atopic dermatitis, psoriasis, eosinophilic gastrointestinal disorder, inflammatory bowel disease*

Innate lymphoid cells (ILCs) are recently identified immune cells that have lymphoid morphology but lack an antigen receptor. Instead of antigen stimulation, ILCs promptly respond to multiple

### Abbreviations used


AD:	Atopic dermatitis
AHR:	Airway hyperresponsiveness
BALF:	Bronchoalveolar lavage fluid
CD:	Crohn disease
cNK:	Classical natural killer
CRT2:	Chemoattractant receptor-homologous molecule expressed on T <sub>H</sub> 2 cells
EGID:	Eosinophilic gastrointestinal disorder
EoE:	Eosinophilic esophagitis
GWAS:	Genome-wide association study
HDM:	House dust mite
IBD:	Inflammatory bowel disease
ILC:	Innate lymphoid cell
ILC1:	Group 1 innate lymphoid cell
ILC2:	Group 2 innate lymphoid cell
ILC3:	Group 3 innate lymphoid cell
KLRG1:	Killer cell lectin-like receptor G1
LTD <sub>4</sub> :	Leukotriene D <sub>4</sub>
LTi:	Lymphoid tissue inducer
NCR:	Natural cytotoxicity receptor
NK:	Natural killer
OVA:	Ovalbumin
PGD <sub>2</sub> :	Prostaglandin D <sub>2</sub>
RORγt:	Retinoic acid-related orphan receptor γt
STAT:	Signal transducer and activator of transcription
T-bet:	T-box transcription factor
Treg:	Regulatory T
TSLP:	Thymic stromal lymphopoietin
UC:	Ulcerative colitis
WAT:	White adipose tissue

From <sup>a</sup>the Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos; <sup>b</sup>the Christine Kühne-Center for Allergy Research and Education, Davos; <sup>c</sup>the Laboratory for Innate Immune Systems and <sup>d</sup>the Laboratory for Immune Cell Systems, RIKEN Center for Integrative Medical Sciences, Yokohama; <sup>e</sup>the Division of Immunobiology, Department of Medical Life Science, Graduate School of Medical Life Science, Yokohama City University; and <sup>f</sup>the Department of Microbiology and Immunology, Keio University School of Medicine, Tokyo.

Disclosure of potential conflict of interest: S. Koyasu had consultant arrangements with Medical and Biological Laboratories. The rest of the authors declare that they have no relevant conflicts of interest.

Received for publication July 19, 2016; revised September 21, 2016; accepted for publication September 22, 2016.

Corresponding author: Shigeo Koyasu, PhD, Laboratory for Immune Cell Systems, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan. E-mail: [shigeo.koyasu@riken.jp](mailto:shigeo.koyasu@riken.jp).

 The CrossMark symbol notifies online readers when updates have been made to the article such as errata or minor corrections

0091-6749

© 2016 The Authors. Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<http://dx.doi.org/10.1016/j.jaci.2016.09.011>

Terms in boldface and italics are defined in the glossary on page 1254.

cell-derived factors, such as cytokines and *eicosanoids*, which are produced by other cells in response to pathogen-associated molecular patterns and *damage-associated molecular patterns*, leading to their designation as innate immune cells. ILCs develop from common lymphoid progenitors in fetal liver and bone marrow and are dependent on *common γ chain* receptor signaling and the transcription factor Id2.<sup>1-3</sup> They have been classified into 3 different subpopulations, namely group 1 innate lymphoid cells (ILC1s), group 2 innate lymphoid cells (ILC2s), and group 3 innate lymphoid cells (ILC3), which resemble T<sub>H</sub> cell subsets, according to their distinct patterns of cytokine production and transcription factors (Fig 1).<sup>4</sup>

ILC1s, which resemble T<sub>H</sub>1 cells, are defined by production of the type 1 cytokine IFN-γ and expression of *T-box transcription factor (T-bet)* and have been implicated in immunity against intracellular pathogens, such as viruses and intracellular

bacteria.<sup>3,5,6</sup> Group 1 ILCs include 2 different populations: classical natural killer (cNK) cells and ILC1s. Both cNK cells and ILC1s produce IFN- $\gamma$  on activation with IL-12, IL-15, and IL-18. The cytotoxic activity of cNK cells is mediated by the production of granzymes and perforin similar to CD8<sup>+</sup> cytotoxic T cells,<sup>7</sup> whereas ILC1s do not produce granzymes or perforin. Although cNK cells develop from Id2<sup>+</sup> progenitors, which have the potential to develop into all ILC subsets, ILC1s develop from a PLZF<sup>high</sup> ILC progenitor that has lost the potential to develop into either cNK cells or lymphoid tissue inducer (LTi) cells, a type of ILC3.

ILC2s, which are phenotypically similar to T<sub>H</sub>2 cells, are defined by the production of the type 2 cytokines IL-4, IL-5, IL-9, and IL-13 and expression of GATA-3.<sup>8-15</sup> Although GATA-3 has been reported to be indispensable for the development of all IL-7 receptor  $\alpha$ <sup>+</sup> ILC subsets (ILC1s, ILC2s, and ILC3s), excluding cNK cells,<sup>16,17</sup> it is a functional requirement for the development of ILC2s. The designation ILC2 collectively refers to natural helper cells and nuocytes (inflammatory ILC2s). Although natural helper cells exist in diverse tissues in the steady state, nuocytes or inflammatory ILC2s are induced in IL-25-mediated inflammation. ILC2s can be activated by various cytokines, such as IL-33, IL-25, *thymic stromal lymphopoietin* (TSLP),<sup>8-11</sup> and IL-1 $\beta$ ,<sup>18,19</sup> as well as eicosanoids, such as prostaglandin D<sub>2</sub> (PGD<sub>2</sub>)<sup>20</sup> and leukotriene D<sub>4</sub> (LTD<sub>4</sub>).<sup>21</sup> ILC2s also produce amphiregulin, a member of the epidermal growth factor family, on activation.<sup>22</sup> Therefore ILC2s have been implicated in immunity against helminths, allergic diseases, and tissue repair.

ILC3s, which resemble T<sub>H</sub>17 cells, are defined by the production of IL-17A and IL-22 and by expression of retinoic acid-related orphan receptor  $\gamma$ t (ROR $\gamma$ t) and have been implicated in immunity against extracellular bacteria and autoimmune diseases.<sup>4</sup> ILC3s

comprise 3 different populations: LTi cells,<sup>23,24</sup> natural cytotoxicity receptor (NCR)<sup>+</sup> ILC3s, and NCR<sup>-</sup> ILC3s. LTi cells express CCR6 and produce IL-17A, IL-17F and IL-22 on activation with IL-1 $\beta$  and IL-23. They have heterogeneous expression of CD4 and are a source of lymphotoxin, which is necessary for the formation of secondary lymphoid tissues.<sup>23,24</sup> NCR<sup>+</sup> ILC3s and NCR<sup>-</sup> ILC3s are distinguished by the expression of NCRs, such as NKp44 in human subjects and NKp46 in mice. Both cell populations produce IL-22 on activation by IL-1 $\beta$  and IL-23,<sup>4,25-31</sup> but NCR<sup>+</sup> ILC3s are strong producers of IFN- $\gamma$ .

Interestingly, ILCs display functional plasticity in response to environmental cues, which enables them to respond promptly to environmental changes. For instance, conversion between ILC1s and ILC3s<sup>30</sup> and ILC1s and ILC2s<sup>18,19,32,33</sup> has been reported in human subjects. In brief, ILC3s differentiate into ILC1s on IL-12 stimulation, followed by T-bet induction, whereas ILC1s revert to ILC3s on stimulation with IL-23, IL-1 $\beta$ , and retinoic acid, which are found in the intestine in patients with Crohn disease (CD).<sup>30</sup> In addition, ILC2s differentiate into ILC1s on IL-12 stimulation, followed by T-bet induction, whereas ILC1s revert to ILC2s on stimulation with IL-4, which is found in the lungs in patients with chronic obstructive pulmonary disease and in nasal tissue in patients with chronic rhinosinusitis with nasal polyps.<sup>18,19,32,33</sup> These findings suggest that ILCs might receive specific tissue-derived signals in response to different types of inflammation and subsequently acquire diverse phenotypes with specialized effector capacities.

## ILCS IN ALLERGIC AIRWAY INFLAMMATION

A recent genome-wide association study (GWAS) showed that genetic polymorphisms in the gene encoding IL-33, which is a

## GLOSSARY

**CHITIN:** Also known as poly-N-acetyl-D-glucosamine, a polysaccharide found in the cell wall of fungi and the integument of insects and crustaceans. Its structure is similar to cellulose. It is used in the pesticide industry as a fertilizer that deters soil nematode growth. Intra-airway administration of chitin induces asthma-like airway inflammation in mice.

**COMMON  $\gamma$  CHAIN:** A signal-transducing chain for multiple type I cytokine receptors (IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21), also known as CD132. Janus kinase 3 (JAK3) is used for common  $\gamma$  chain signaling. Common  $\gamma$  chain mutations are the cause of X-linked severe combined immunodeficiency (X-SCID). Autosomal recessive mutations in the JAK3 gene result in a similar phenotype as X-SCID.

**DAMAGE-ASSOCIATED MOLECULAR PATTERNS (DAMPs):** Molecules produced by cell damage caused by infection, toxins, burns, trauma, or hypoperfusion. Alarmins are a specific DAMP produced to enhance the innate immune response to infection. Other examples include heat shock protein and monosodium urate.

**EICOSANOIDS:** Arachidonic acid-derived compounds that act as inflammatory mediators. Examples include prostaglandins and leukotrienes. The term is based on the Greek word *eikosi*, which means 20 (the number of carbon atoms in arachidonic acid).

**IMIQUIMOD:** A Toll-like receptor 7 ligand used as a pharmaceutical topical product for the treatment of condyloma acuminata, actinic keratosis, and superficial basal cell carcinoma. Topical application of imiquimod induces psoriasis-like skin inflammation in mice.

**LAMINA PROPRIA:** A layer of the mucus membrane that lies beneath the epithelium and contains connective tissue, lymphocytes, mast cells, and plasma cells.


**PAPAIN:** A cysteine protease present in papaya fruit. One of its main uses in the food industry is as a meat-tenderizing agent. Intra-airway administration of papain induces asthma-like airway inflammation in mice.

**RECOMBINATION-ACTIVATING GENE (*Rag*):** A lymphoid-specific gene that produces protein critical for V(D)J recombination. *Rag* stands for recombination-activating gene and includes *Rag1* and *Rag2* proteins. *Rag1* or *Rag2* deficiency leads to the deficiency of T, B, and NKT cells and results in severe combined immunodeficiency.

**STEROID-RESISTANT ASTHMA:** Often defined in the literature as an adult asthmatic patient whose FEV<sub>1</sub> percent predicted does not improve by 10% to 12% after a course of 20 mg of prednisone twice daily for 7 days.

**T-box TRANSCRIPTION FACTOR (T-bet):** A member of the T-box family of transcription factors that is encoded by the gene *Tbx21* and activated by IFN- $\gamma$ . Its induction in CD4<sup>+</sup> T cells then stimulates additional IFN- $\gamma$  production, resulting in a positive amplification loop for T<sub>H</sub>1 differentiation.

**THYMIC STROMAL LYMPHOPHOETIN:** A cytokine secreted by gastrointestinal, skin, and lung epithelial cells that promotes a T<sub>H</sub>2 differential through its effect on tissue dendritic cells and ILC2s. Its name arises from its initial identification in thymic stromal cells.




**FIG 1.** Effector subsets of ILCs. ILCs are divided into 3 different subsets based on their different transcription factors and cytokine expression. ILC1s produce type 1 cytokines, such as IFN- $\gamma$ , on stimulation with IL-12, IL-15, and IL-18 and express T-bet. These cells are involved in immune responses against intracellular pathogens and act in chronic inflammation. Group 2 ILCs produce type 2 cytokines, such as IL-4, IL-5, IL-9, and IL-13, on stimulation with epithelium-derived cytokines, such as IL-33, IL-25, TSLP, eicosanoids, and IL-1 $\beta$ . These cells are involved in immune responses against helminths or fungi and induce allergic inflammation. ILC3s produce type 3 cytokines, such as IL-17A and IL-22, in response to IL-1 $\beta$  and IL-23 stimulation. These cells play an important role in defense against extracellular pathogens and chronic inflammation. *Areg*, Amphiregulin; *Eomes*, eomesodermin.

major activator of ILC2s, and its receptor IL-1RL1 (ST2) are strongly linked to asthma development.<sup>34,35</sup> Although the genes that encode IL-33 and IL-1RL1 are located on different chromosomes, these 2 genes have consistently been identified in many reports, strongly suggesting that the IL-33–IL-1RL1 pathway and ILC2s are involved in asthma pathogenesis.<sup>36</sup> Indeed, in human subjects a recent study showed that the frequency of ILC2s, as defined by the expression of chemoattractant receptor-homologous molecule expressed on T<sub>H</sub>2 cells (CRTH2), IL-7 receptor in peripheral blood<sup>37–39</sup> and bronchoalveolar lavage fluid (BALF),<sup>39–41</sup> and IL-33 levels in BALF,<sup>40</sup> were increased in asthmatic patients compared with control subjects. Moreover, the frequency of ILC2s in peripheral blood<sup>38</sup> and BALF,<sup>40</sup> as well as IL-33 levels in BALF,<sup>40</sup> were negatively correlated with airway function, indicating that ILC2s and IL-33 are involved in asthma pathogenesis.

A recent study showed that ILC2s are involved in eosinophilic asthma-like airway inflammation, including eosinophil

infiltration, airway hyperresponsiveness (AHR), and mucus production, even in the absence of acquired immunity, such as T cells, B cells, and IgE antibody.<sup>42–44</sup> In addition, in mice inhalation of epithelium-derived cytokines, such as IL-33 and IL-25, induces eosinophilic asthma-like airway inflammation accompanied by AHR, even in *recombination-activating gene (Rag)*<sup>−/−</sup> mice, which lack T cells, B cells, and natural killer (NK) T cells.<sup>42–46</sup> In contrast, IL-33-induced airway inflammation was diminished in *Rag2*<sup>−/−</sup>*Il2rg*<sup>−/−</sup> mice, which lack all lymphocytes, including ILCs, but was restored to some extent by engraftment of ILC2s.<sup>42</sup> These findings clearly show that ILC2s induce eosinophilic airway inflammation independent of acquired immunity on activation by epithelium-derived cytokines, such as IL-33 and IL-25 (Fig 2). It has been shown recently that many asthma-related antigens, such as protease allergens, fungal extracts, and viral infection, trigger IL-33 and IL-25 production from epithelial cells and various immune cells and induce eosinophilic asthma-like airway inflammation through activation of lung ILC2s



**FIG 2.** Roles of ILCs in allergic and nonallergic airway inflammation. In patients with allergic airway inflammation, protease allergens, such as HDM, and fungal and viral infection induce IL-33, IL-25, and TSLP from epithelial cells and macrophages. These cytokines induce production of type 2 cytokines, such as IL-5 and IL-13, from ILC2s, which leads to eosinophilic inflammation, even in the absence of acquired immunity. PGD<sub>2</sub> and LTD<sub>4</sub> from mast cells also induce type 2 cytokines from ILC2s, and basophil-derived IL-4 and inducible costimulator (ICOS)/ICOS ligand interaction on ILC2s are important for ILC2 activation. IL-5 from ILC2s promotes accumulation of eosinophils, and IL-13 induces goblet cell hyperplasia and accumulation of alveolar macrophages. ILC2s also activate T<sub>H</sub>2 cells through OX40 ligand (OX40L) and MHC class II. IFN-α/β, IFN-γ, IL-27, prostaglandin I<sub>2</sub> (PGI<sub>2</sub>), and Treg cells suppress ILC2 activation and attenuate ILC2-mediated airway inflammation. In patients with nonallergic airway inflammation, such as obesity-associated asthma, IL-1β derived from macrophages induces IL-17A production by ILC3s, which results in induction of neutrophils and AHR. Macrophage-derived IL-1β can also activate ILC2s and enhance airway inflammation. Areg, Amphiregulin.

independent of acquired immunity.<sup>42,44,47-49</sup> Furthermore, many allergens, such as house dust mites (HDMs), cockroaches, fungi, and pollens were reported to possess protease activities.<sup>50</sup> Although protease activities of allergens destroy epithelial cells, which enables the allergen to cross the tissue barrier, resulting in allergen sensitization, these protease activities also induce IL-33 release from epithelial cells, which induces asthma-like airway inflammation through ILC2 activation. In fact, *papain*, which is homologous to the HDM allergen Der 1 and used in the food industry, is known to be a cause of occupational asthma.<sup>51</sup> Administration of papain to mice induces IL-33 secretion from lung epithelial cells and induces eosinophilic asthma-like airway inflammation through activation of lung ILC2s by IL-33.<sup>42,44,47,52</sup> Under these conditions, it was shown that basophil-derived IL-4<sup>53</sup> and inducible costimulator/inducible costimulator ligand interaction on ILC2s<sup>54</sup> play important roles in the activation of lung ILC2s. However, mast cells activated by IL-33 play a protective role by expanding regulatory T (Treg) cells that suppress ILC2 activation (Fig 2).<sup>42,55</sup>

Administration of *Alternaria* species, a major fungal aeroallergen, also induces eosinophilic airway inflammation in mice

through activation of lung ILC2s by IL-33 and IL-25.<sup>21,47,48</sup> In agreement with this, it was shown that IL-33 expression was increased in sinonasal epithelial cells from patients with chronic rhinosinusitis with nasal polyps on stimulation with fungal extracts from *Alternaria alternata*, *Aspergillus fumigatus*, and *Cladosporium herbarum*.<sup>56</sup> *Chitin*, a key structural component of fungi, HDM, and helminths, is also known to induce eosinophilic airway inflammation independent of acquired immunity.<sup>57</sup> Recently, it was shown that ILC2s activated by IL-33, IL-25, and TSLP are essential for the induction of chitin-induced eosinophilic airway inflammation through type 2 cytokine production.<sup>58</sup>

Viral infection also induces asthma-like airway inflammation through activation of ILC2s independent of acquired immunity. For example, it was shown that H3N1 influenza A virus infection induced AHR through activation of lung ILC2s by alveolar macrophage-derived IL-33.<sup>49</sup> In addition, respiratory syncytial virus infection induced AHR and mucus accumulation through activation of lung ILC2s by epithelium-derived TSLP but not IL-33.<sup>59,60</sup> Rhinovirus infection of neonatal mice induced long-term mucus metaplasia and AHR through activation of lung ILC2s by epithelium-derived IL-25.<sup>61</sup> Interestingly, unlike



neonatal mice, IL-25 production from epithelial cells and ILC2 accumulation were not observed in adult mice infected by rhinovirus,<sup>61</sup> suggesting that early-life rhinovirus infection might contribute to the development of asthma later through activation of lung ILC2s, which results in chronic mucus metaplasia and AHR. In support of this notion, a recent epidemiologic study showed that wheezing episodes on viral respiratory tract infections, such as rhinovirus and respiratory syncytial virus in early life, are strongly associated with asthma development.<sup>62</sup> Taken together, activation of ILC2s by IL-33, IL-25, and TSLP induced by allergens and viruses is crucial for initiation of asthma-like airway inflammation development before antigen sensitization.

ILC2s contribute not only to the initiation of innate allergic airway inflammation but also to the enhancement of allergic inflammation by interacting with other immune cells. The role of ILC2s in acquired immunity was analyzed in mouse models that require a sensitization phase before the challenge phase, such as HDM, ovalbumin (OVA), and papain exposure models. In an HDM model and papain exposure model, eosinophilic airway inflammation was attenuated in mice that were lethally irradiated and transplanted with bone marrow cells from *Rora*<sup>sg/sg</sup> mice. Because *Rora* is critical for ILC2 differentiation, these mice lack ILC2s (ILC2-deficient mice). T-cell responses to antigen were significantly decreased in ILC2-deficient mice compared with control mice.<sup>63,64</sup> ILC2s have been reported to have a direct effect on T<sub>H</sub>2 cell activation through OX40 ligand<sup>65</sup> and MHC class II<sup>66</sup> on ILC2s *in vitro*. In addition, ILC2s have also been reported to have an indirect effect on memory T<sub>H</sub>2 cell recruitment through induction of CCL17 from dendritic cells.<sup>67</sup> These mechanisms might be involved in the activation of acquired immunity (Fig 2). In contrast, in an OVA model eosinophilic airway inflammation was comparable between ILC2-deficient mice and control mice.<sup>63</sup> Whereas HDM and papain exposure model mice were sensitized by intranasal antigen administration, OVA model mice were sensitized by systemic antigen administration. Therefore it is possible that ILC2s have crucial roles in the initiation of local immune responses to antigen but not in the initiation of systemic immune responses.

In human lungs ILC2s are located in proximity to mast cells,<sup>68</sup> suggesting that ILC2s can also interact with mast cells. CRTH2, a distinctive marker on human ILC2s, is a receptor for PGD<sub>2</sub> that is produced by activated mast cells. It was shown that PGD<sub>2</sub> induces migration of human ILC2s through CRTH2<sup>20,69</sup> and induces type 2 cytokine production from human ILC2s.<sup>20,68</sup> Another lipid mediator, LTD<sub>4</sub>, derived from mast cells was also shown to activate ILC2s in mice. LTD<sub>4</sub> induces type 2 cytokine production from mouse ILC2s and accumulation of lung ILC2s in *Alternaria* species-induced eosinophilic inflammation.<sup>21</sup> PGD<sub>2</sub> and LTD<sub>4</sub> are released by mast cells on degranulation induced by IgE cross-linking.<sup>70,71</sup> Therefore these findings suggest that mast cells activated on IgE cross-linking contributes to enhancement of allergic airway inflammation through recruitment and activation of ILC2s. In contrast, IL-33 induces IL-2 production from mast cells that promotes Treg cell expansion<sup>42</sup> but not degranulation of mast cells.<sup>72,73</sup> Therefore mast cells activated by IL-33 contribute to the regulation of allergic airway inflammation through suppression of ILC2s by Treg cells (Fig 2).

Recent studies suggest that ILC2s also have important roles in steroid-resistant severe allergic airway inflammation and persistence of chronic airway inflammation.<sup>39,43</sup> In human subjects it was shown that the number of total ILC2s and type 2 cytokine-

producing ILC2s in peripheral blood and sputum were significantly increased in patients with systemic steroid-dependent severe eosinophilic asthma compared with those with mild asthma. In contrast, the number of type 2 cytokine-producing CD4<sup>+</sup> T cells did not differ significantly between these 2 patient groups.<sup>39</sup> These findings suggest that ILC2s, but not CD4<sup>+</sup> T cells, play a significant role in **steroid-resistant asthma** and promote the development of airway inflammation in patients with systemic steroid-dependent severe eosinophilic asthma. Likewise, corticosteroids suppressed T<sub>H</sub>2 cells, but not ILC2s, in an OVA-induced asthma model when a low dose of IL-33 was coadministered with OVA, which results in corticosteroid-resistant airway inflammation. In this model TSLP was shown to be a key molecule that induces corticosteroid resistance of ILC2s by controlling phosphorylation of signal transducer and activator of transcription (STAT) 5 and expression of Bcl-xL. Furthermore, STAT5 inhibitors abrogated steroid resistance in this model.<sup>43</sup>

It is known that symptoms persist in most patients with occupational asthma, even years after removal of exposure to the occupational antigen.<sup>74</sup> However, little is known about the mechanisms regulating these persistent symptoms. It was recently shown that ILC2s, but not T cells, play a pivotal role in the persistence of asthma. Three allergens (ie, HDM, ragweed, and *Aspergillus* species) induced persistent allergic inflammation in mice, which exhibited persistent inflammation and AHR for more than 6 months after cessation of allergen exposure. In these models depletion of T cells reduced sustained airway inflammation to some extent but did not eliminate sustained AHR. In contrast, depletion of ILC2s completely eliminated sustained airway inflammation and AHR, indicating the importance of ILC2s in the persistence of chronic asthma.<sup>40</sup>

Taken together, ILC2s play a pivotal role in the initiation, enhancement, and steroid resistance of allergic airway inflammation. Consequently, regulation of ILC2 activation is likely an important target to control asthma. Recently, IFN- $\alpha/\beta$ ,<sup>75,76</sup> IFN- $\gamma$ ,<sup>75-77</sup> IL-27,<sup>75,76,78</sup> Treg cells,<sup>42</sup> and prostaglandin I<sub>2</sub>,<sup>79</sup> were shown to suppress ILC2 activation *in vitro* and *in vivo* (Fig 2). These findings open a new window for the prevention and treatment of asthma.

## ILCs IN NONALLERGIC AIRWAY INFLAMMATION

Although asthma has been considered a single disease for years, recent studies indicate that asthma is actually a heterogeneous disease composed of different phenotypes, such as allergic asthma, including early-onset allergic asthma and late-onset persistent eosinophilic asthma, and nonallergic asthma, including neutrophilic asthma and obesity-related asthma.<sup>80</sup> As mentioned before, evidence has accumulated that ILC2s are involved in allergic-type asthma. However, the role of ILCs in nonallergic asthma remains unclear. Obesity has been suggested to play a substantial role in the development of asthma.<sup>81</sup> Recent findings suggest that ILCs are also involved in obesity-associated allergic airway inflammation. The numbers of ILC3s in the lungs of high-fat diet-fed obese mice was increased compared with those in standard diet-fed mice. In addition, AHR was induced in high-fat diet-fed obese mice in the absence of an antigen challenge. This obesity-induced AHR is independent of acquired immunity but dependent on IL-17A. Obesity-induced AHR was ameliorated in *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice, which lack all ILCs in addition to T cells, B cells, and NKT cells, and adoptive transfer of ILC3s

to *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice restored obese-induced AHR, suggesting that IL-17A-producing ILC3s play a critical role in obesity-induced AHR. Under these conditions, IL-1 $\beta$  produced by macrophages on activation of the nucleotide-binding domain, leucine-rich repeat/pyrin domain-containing 3 pathway induces IL-17A production from ILC3s (Fig 2).<sup>82</sup>

It has recently been shown that obesity contributes to airway inflammation not only through activation of ILC3s but also through ILC2 activation. ILC2 numbers in the lungs of high-fat diet-fed obese mice were also increased compared with those in low-fat diet-fed mice without antigen challenge. On HDM administration, airway inflammation and AHR accompanied by accumulation of ILC2s and ILC3s were exacerbated in high-fat diet-fed obese mice compared with mice fed a standard diet, suggesting that activation of both ILC2s and ILC3s plays an important role in obesity-associated airway inflammation on HDM challenge.<sup>83</sup> Although the mechanisms for ILC2 activation in high-fat diet-fed obese mice have not been clarified, a recent study showed that IL-1 $\beta$  also activates ILC2s to produce type 2 cytokines,<sup>18,19</sup> suggesting that IL-1 $\beta$  signaling is a possible mechanism underlying obesity-induced ILC2 activation (Fig 2).

Interestingly, unlike in the lungs, it was shown that ILC2 responses in white adipose tissue (WAT) were dysregulated in obese patients. Because the IL-33/ILC2 axis regulates metabolic homeostasis by eliciting beiging of WAT, dysregulation of ILC2s in WAT might be one possible mechanism of obesity.<sup>84</sup> However, the mechanisms that induce dysregulation of ILC2 responses in WAT remain unclear.

Of note, ILC2s promote beneficial tissue repair responses in the lung after acute epithelial damage, such as viral infection. It has been reported that depletion of ILC2s during H1N1 influenza infection impaired lung function and epithelial integrity, which were restored after adoptive transfer of lung ILC2s or administration of amphiregulin, suggesting that ILC2s contribute to lung tissue repair through amphiregulin production (Fig 2).<sup>22</sup> However, as mentioned before, by using a different H3N1 influenza virus, another article showed that ILC2s could be pathogenic during influenza virus infection through IL-13 production.<sup>49</sup> These findings suggest that the roles of ILC2s in the lungs during viral respiratory tract infection might be different depending on the level of tissue damage, types of virus, and environmental factors. Further studies are needed to fully characterize the different roles of ILC2s in the lungs during viral respiratory tract infection.


## ILCS IN ALLERGIC SKIN INFLAMMATION

Atopic dermatitis (AD) is the most common chronic and relapsing inflammatory skin disease characterized by type 2-skewed inflammation and epithelial barrier dysfunction.<sup>85</sup> Expression of TSLP,<sup>86</sup> IL-25,<sup>87</sup> and IL-33,<sup>88</sup> which activate ILC2s, was upregulated in lesional skin from patients with AD compared with healthy skin, suggesting that ILC2s play a crucial role in the pathogenesis of AD through activation by these cytokines. Indeed, it has been shown that ILC2s are resident in healthy skin tissue in both human subjects and mice,<sup>89,90</sup> and the frequency of ILC2s was higher in skin tissue from lesional skin of human patients with AD compared with that in healthy subjects.<sup>89,90</sup> The responsiveness of skin ILC2s to epithelium-derived cytokines is still controversial. Salimi et al<sup>90</sup> reported that ILC2s isolated from healthy skin produced type 2 cytokines

on stimulation by IL-33 but not IL-25 and TSLP. Conversely, Teunissen et al<sup>91</sup> reported that ILC2s from healthy skin samples produced type 2 cytokines on TSLP, but not IL-33 and IL-25, stimulation. However, IL-17RB (IL-25 receptor), ST2 (IL-33 receptor), TSLP receptor, and CCR2 were highly upregulated on ILC2s from human AD skin lesions compared with ILC2s from healthy skin,<sup>90</sup> suggesting that ILC2s in AD skin lesions are highly activated and more responsive to stimuli, such as IL-25, IL-33, TSLP, and PGD<sub>2</sub> (Fig 3). The role of ILC2s in allergic skin inflammation has also been investigated by using different mouse models. Overexpression of IL-33<sup>92</sup> and TSLP<sup>93,94</sup> in the skin spontaneously induces AD-like skin inflammation accompanied by itching, eosinophil infiltration, and thickening of the epidermis. In particular, numbers of Lin<sup>-</sup>ST2<sup>+</sup>Sca-1<sup>+</sup> skin ILC2s were increased in lesional skin from transgenic mice with skin-specific expression of IL-33, indicating involvement of the IL-33-ILC2 axis in this model.<sup>92</sup> Topical treatment with a vitamin D analog, calcipotriol, was also shown to induce AD-like skin inflammation accompanied by eosinophil infiltration, epidermal hyperplasia, and increased type 2 cytokine production.<sup>89</sup> Similarly, a vitamin D analog also induced AD-like skin inflammation in *Rag1*<sup>-/-</sup> mice, which lack acquired immunity, but depletion of ILCs by an anti-CD25 antibody attenuated skin inflammation, suggesting that ILCs, but not T cells, B cells, and NKT cells, are critical in this model.<sup>89</sup> It is unknown whether similar mechanisms mediate skin inflammation in the above-mentioned mouse models.

Furthermore, the involvement of epithelium-derived cytokines, such as IL-33 and TSLP, in these models is controversial. Kim et al<sup>89</sup> demonstrated that skin inflammation was attenuated in *Tslpr*<sup>-/-</sup> mice, but not in *Il33*<sup>-/-</sup> mice, compared with wild-type mice on a C57BL/6 background. In contrast, Salimi et al<sup>90</sup> demonstrated that skin inflammation was attenuated in *Il17rb*<sup>-/-</sup> mice and *St2*<sup>-/-</sup> mice, but not *Tslpr*<sup>-/-</sup> mice, on a BALB/c background. These differences might be due to differences in genetic backgrounds and environmental factors, including microbiota.

Interactions of ILC2s with other cell types have also been shown to be critical in the pathogenesis of allergic skin inflammation. Basophils, which are usually undetectable in healthy control skin, were enriched in the dermis of human AD lesions in close proximity to ILC2s,<sup>95</sup> suggesting that the interaction between ILC2s and basophils is involved in the pathogenesis of AD. Indeed, depletion of basophils attenuated vitamin D analog-induced AD-like skin inflammation in mice. Moreover, conditional depletion of IL-4 in basophils also attenuated vitamin D analog-induced AD-like skin inflammation and accumulation of ILC2s in the skin.<sup>95</sup> Therefore it is likely that basophils play a critical role in the initiation of allergic skin inflammation through activation of skin ILC2s by means of production of IL-4 (Fig 3). Mast cells also accumulated in the lesional skin in human and mouse AD models,<sup>96</sup> suggesting that the interaction between mast cells and ILC2s is also involved in the pathogenesis of AD. Indeed, it has been shown that PGD<sub>2</sub> promotes the migration of skin ILC2s and induces type 2 cytokine production from skin ILC2s. Moreover, the supernatant of IgE/anti-IgE-activated mast cells also induces migration and cytokine production by human skin ILC2s.<sup>20</sup> Because mast cells produce PGD<sub>2</sub> by means of activation through Fc $\epsilon$ RI, they likely play a role in migration and cytokine production by ILC2s through the effects of PGD<sub>2</sub> (Fig 3). Killer cell lectin-like receptor G1 (KLRG1), which is known



**FIG 3.** Role of ILCs in allergic and nonallergic skin inflammation. In patients with allergic skin inflammation, such as AD, epithelial cell–derived cytokines, such as IL-33, IL-25, and TSLP, induce type 2 cytokine production from ILC2s. ILC2s from AD lesional skin express IL-17RB (IL-25 receptor), ST2 (IL-33 receptor), TSLP receptor, and CRTH2. Basophils and mast cells accumulate in the lesional skin of patients with AD. Similar to what is observed in patients with allergic airway inflammation, basophil-derived IL-4 is important for ILC2 activation. Mast cell–derived PGD<sub>2</sub> induces type 2 cytokine production from ILC2s. E-cadherin expressed on keratinocytes and Langerhans cells suppresses ILC2 activation by means of ligation to KLRG1 on ILC2s. Decreased expression of E-cadherin in keratinocytes from patients with AD results in persistent activation of ILC2s. In patients with nonallergic skin inflammation, such as psoriasis, IL-22 from NCR<sup>+</sup> ILC3s induces skin thickening, and IL-17A from ILC3s induces neutrophil infiltration. ILC3s accumulate in wound sites through Notch1 and contribute to wound healing through IL-17F, IL-17A, and IL-22.

as an inhibitory receptor on NK cells, is also a known marker of ILC2s in mice.<sup>97</sup> KLRG1 binds to the cell adhesion molecule E-cadherin, which is expressed on keratinocytes and Langerhans cells.<sup>98</sup> KLRG1 was highly expressed on skin ILC2s from patients with AD compared with skin ILC2s from healthy control subjects, and TSLP and IL-33, but not IL-25, upregulated KLRG1 expression on cultured ILC2s. In addition, E-cadherin ligation to KLRG1 on ILC2s suppressed cytokine production from skin ILC2s,<sup>90</sup> suggesting that the KLRG1–E-cadherin interaction plays an important role in the negative feedback mechanisms in activated skin ILC2s (Fig 3). Interestingly, E-cadherin expression was decreased in keratinocytes from AD lesional skin compared with that seen in normal skin.<sup>99</sup> In addition, E-cadherin was downregulated in keratinocytes on short hairpin RNA knockdown of the filaggrin gene that has crucial roles in skin barrier function.<sup>90</sup> These findings suggest that low E-cadherin expression on keratinocytes in patients with AD leads to persistent activation of ILC2s in AD skin lesions.

Taken together, ILC2s are likely involved in the induction of allergic skin inflammation through activation by epithelium–derived cytokines, such as IL-33, IL-25, and TSLP, and interaction with other immune cells.

### ILCs in Nonallergic Skin Inflammation

Psoriasis is a common chronic skin inflammatory disease characterized by epidermal thickness and scaly plaques that affects approximately 2% of the population.<sup>100</sup> Previously, T cells and dendritic cells were considered to play key roles in the pathogenesis of psoriasis through production of cytokines, such as IL-17A, IL-17F, and IL-22. In brief, IL-17A and IL-17F contribute to disease pathogenesis by inducing neutrophilic inflammation, and IL-22 contributes to epidermal hyperplasia (Fig 3).<sup>101</sup> However, recent studies have shown that ILCs, particularly ILC3s that produce IL-17A and IL-22, also play substantial roles in the pathogenesis of psoriasis. The frequency of NCR<sup>+</sup> ILC3s, but not ILC1s or ILC2s, was increased in peripheral blood from patients with psoriasis compared with healthy control subjects.<sup>91,102</sup> Moreover, the frequency of NCR<sup>+</sup> ILC3s was increased in lesional<sup>91</sup> and nonlesional<sup>103</sup> skin from patients with psoriasis compared with healthy skin. Importantly, the frequency of NCR<sup>+</sup> ILC3s in the skin correlated with disease severity determined by the Psoriasis Area and Severity Index.<sup>91</sup> In addition, the frequency of NCR<sup>+</sup> ILC3s in peripheral blood was decreased on anti-TNF antibody treatment and was closely associated with the reduction of disease severity determined by using the Psoriasis



Area and Severity Index.<sup>102</sup> ILC3s from lesional skin of patients with psoriasis can produce IL-22 and, to a lesser extent, IL-17A,<sup>91,102</sup> suggesting that ILC3s have a pathogenic role in psoriasis through IL-22 and, to a lesser extent, IL-17A production.

In mice topical application of *imiquimod* induces skin inflammation accompanied by acanthosis and plaque formation that largely resembles the disease phenotype of human psoriasis. In this model  $\gamma\delta$  T cells and ILC3s, but not  $T_H17$  cells, were the main source of these cytokines. Indeed, imiquimod-induced skin inflammation was induced in *Rag1*<sup>-/-</sup> mice, which lack acquired immunity, but was completely diminished in *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice, which lack ILCs, and *Rorgt*<sup>-/-</sup> mice, which lack  $T_H17$  cells and ILC3s.<sup>104</sup> Taken together, these findings suggest that ILC3s have critical roles as effector cells in the development of psoriasis-like skin inflammation in mice through production of IL-17A, IL-17F, and IL-22 (Fig 3).

ILC3s have also been shown to play a substantial role in skin repair. It was shown that  $ROR\gamma t^+$  ILC3s were recruited to wound sites through epidermal Notch1 signaling in mice. Similarly  $ROR\gamma t^+$  ILC3s infiltrated human skin after wounding by using a punch biopsy. In addition, wound healing was delayed in mice deficient for  $ROR\gamma t^+$  ILC3s accompanied by delayed epidermal proliferation and macrophage recruitment.<sup>105</sup> Interestingly, the majority of mouse ILC3s that accumulated in the wound site expressed IL-17F but lower amounts of IL-17A and IL-22. Likewise, human ILC3s accumulating in the wound site expressed IL-17F and IL-17A.<sup>105</sup> These findings suggest that ILC3s contribute to repair of skin tissue through production of IL-17F and, to a lesser extent, IL-17A and IL-22 (Fig 3).

## ILCs IN ALLERGIC GASTROINTESTINAL INFLAMMATION

Primary eosinophilic gastrointestinal disorders (EGIDs), including eosinophilic esophagitis (EoE), eosinophilic gastritis, eosinophilic gastroenteritis, and eosinophilic colitis, are disorders that exhibit eosinophil-rich inflammation in the gastrointestinal tract in the absence of known causes for eosinophilia such as parasite infection and drug reaction.<sup>106,107</sup> Although food-specific IgE antibodies were detected in the serum of patients with EGIDs, typical symptoms observed in patients with IgE-dependent food allergy, such as anaphylaxis, were not induced in patients with EGIDs. Therefore EGIDs were considered to be independent of IgE antibodies but dependent on IL-5<sup>+</sup>IL-13<sup>+</sup>  $T_H2$  cell activation by antigens.<sup>106</sup> However, recent findings suggest the involvement of ILC2s in the pathogenesis of EGIDs.

EoE is a chronic, antigen-mediated disease characterized by eosinophil infiltration, esophageal remodeling, subepithelial fibrosis, and smooth muscle dysmotility. Although type 2 cytokines, such as IL-5 and IL-13, were shown to be involved in its pathogenesis,<sup>108,109</sup> the cellular sources of these cytokines were not well clarified. Results from GWASs suggest that the *TSLP* gene is associated with EoE.<sup>110</sup> Indeed, TSLP expression was increased in esophageal biopsy specimens from patients with active EoE compared with control subjects.<sup>110,111</sup> In addition, expression of IL-33 and IL-25 was increased in the esophagus, and the frequency of ILC2s was increased in esophageal biopsy specimens from patients with active EoE compared with those with inactive EoE and control subjects. Importantly, the frequency of ILC2s correlated with the number of eosinophils in esophageal biopsy samples from patients with EoE.<sup>112</sup> These

findings strongly suggest that ILC2s have a pathogenic role in patients with EoE through type 2 cytokine production induced by TSLP, IL-33, and IL-25 (Fig 4).

The involvement of ILCs in patients with other types of EGIDs remains unclear. However, it was shown that TSLP and IL-33 expression in sigmoid colon biopsy specimens was increased in patients with eosinophilic gastroenteritis,<sup>113</sup> suggesting that ILC2s are also involved in other types of EGIDs.

## ILCs IN NONALLERGIC GASTROINTESTINAL INFLAMMATION


Inflammatory bowel diseases (IBDs), such as CD and ulcerative colitis (UC), are a group of chronic disorders of the gastrointestinal tract characterized by intestinal inflammation and epithelial injury.<sup>114</sup> GWASs have identified several IBD susceptibility loci that contain genes encoding cytokines and cytokine receptor signaling. These loci contain genes related to type 1 immune responses, such as those encoding IFN- $\gamma$  and IL-12 receptor and its intracellular signaling components, such as STAT1 and STAT4, as well as genes related to type 3 immune responses, such as those encoding IL-23 receptor and its intracellular signaling components, such as STAT3.<sup>115</sup> These findings suggest that type 1 and type 3 cytokines are involved in the pathogenesis of IBD. Indeed, various studies showed that production of the type 1 cytokine IFN- $\gamma$  by *lamina propria*  $T_H1$  cells was increased in patients with CD compared with healthy control subjects but not in patients with UC.<sup>116,117</sup> In addition, production of type 3 cytokines, such as IL-17A and IL-17F, by lamina propria  $T_H17$  cells was increased in patients with CD, as well as in patients with UC.<sup>117,118</sup> ILCs also produce type 1 and type 3 cytokines similar to T cells, but their involvement in IBD *in vivo* is largely unknown.

The frequency of ILC1s in noninflamed intestine was reported to be approximately 10% of total ILCs but increased to 40% in inflamed intestines of patients with CD.<sup>5,30</sup> In human subjects it was shown that there are 2 distinctive subtypes of ILC1s, namely CD127<sup>+</sup> ILC1s located in the lamina propria<sup>5,30</sup> and CD103<sup>+</sup>CD127<sup>-</sup> ILC1s located in the epithelium.<sup>6</sup> Both populations produce IFN- $\gamma$  on IL-12 stimulation in synergy with IL-15 and IL-18. Whereas the frequencies of CD127<sup>+</sup> ILC1s and CD103<sup>+</sup>CD127<sup>-</sup> ILC1s were comparable in noninflamed intestinal tissue, the frequency of CD127<sup>+</sup> ILC1s, but not CD103<sup>+</sup>CD127<sup>-</sup> ILC1s, was significantly increased in inflamed intestinal tissues from patients with CD.<sup>30</sup> These findings suggest that ILC1s play a role in the pathogenesis of CD through IFN- $\gamma$  production (Fig 4).

ILC2s were also detected in intestinal tissue from patients with CD. Interestingly, unlike ILC2s in peripheral blood, ILC2s in intestinal tissue from patients with CD produce IFN- $\gamma$  in addition to IL-13.<sup>32</sup> These findings suggest that ILC2s in intestinal tissue from patients with CD might have received environmental cues, which enable them to convert to ILC1s.

ILC3s comprise the majority of ILCs (NCR<sup>+</sup> ILC3s, 70%; NCR<sup>-</sup> ILC3s, 15%) in noninflamed human intestine.<sup>5,30</sup> Although the frequency of NCR<sup>-</sup> ILC3s in inflamed intestine from patients with CD was comparable with that in noninflamed control intestinal tissue, the frequency of NCR<sup>+</sup> ILC3s was dramatically (70%  $\rightarrow$  20%) decreased in inflamed intestine from patients with CD.<sup>5,119</sup> Intestinal NCR<sup>+</sup> ILC3s produce IL-22 in response to IL-23 in synergy with IL-1 $\beta$ .<sup>5,120</sup> In addition, intestinal CD14<sup>+</sup> macrophages promote IL-22 production from NCR<sup>+</sup> ILC3s through IL-23 and direct interaction (Fig 4).<sup>31</sup>





**FIG 4.** Roles of ILCs in allergic and nonallergic gastrointestinal inflammation. In patients with allergic gastrointestinal inflammation, such as EoE, epithelium-derived cytokines, such as IL-33, IL-25, and TSLP, induce type 2 cytokine production from ILC2s and induce eosinophil infiltration. In patients with nonallergic gastrointestinal inflammation, such as IBD, ILC1s contribute to disease pathogenesis through IFN- $\gamma$  production. Macrophage-derived IL-23 induces IL-22 production from NCR<sup>+</sup> ILC3s, which is important for intestinal barrier function, wound healing, and antimicrobial production. Decreased frequency of NCR<sup>+</sup> ILC3s might unfavorably affect the pathogenesis of CD. NCR<sup>-</sup> ILC3s produce IL-17A together with IL-22. However, the role of IL-17A in human IBD is not clear. ILC3s also have protective roles in the intestine. ILC3s induce cell death in T<sub>H</sub>17 cells through MHC class II. In addition, ILC3s produce GM-CSF, which activates macrophages to generate Treg cells by retinoic acid and TGF- $\beta$ .

The receptor for IL-22 is mainly expressed on tissue cells, such as epithelial cells, and IL-22 mediates intestinal epithelial barrier maintenance and wound healing<sup>31,121,122</sup> and provides functional barrier support by inducing antimicrobials, such as Reg family molecules and S110 proteins.<sup>123</sup> These findings suggest that IL-22 is important for protection against the bacterial infection and tissue injury that occur in patients with IBD. Indeed, in mice ILC3s were an essential source of IL-22 that protected against dextran sodium sulfate–induced colitis, in which the epithelial barrier was disrupted.<sup>124,125</sup> Thus NCR<sup>+</sup> ILC3s might play a protective role in the pathogenesis of IBD through IL-22 production in humans, and the decrease of IL-22–producing NCR<sup>+</sup> ILC3s is unfavorable for patients with IBD (Fig 4). Intriguingly, however, IL-22 was also pathogenic in certain mouse models that mimic human IBD.<sup>126</sup> Further investigation is needed to clarify the roles of IL-22–producing NCR<sup>+</sup> ILC3s in patients with IBD.

Another subset of ILC3s, NCR<sup>-</sup> ILC3s, have also been described in the human intestine. Human intestinal NCR<sup>-</sup> ILC3s from patients with CD produce IL-17A and IL-17F and, to a lesser extent, IL-22 on IL-23 stimulation.<sup>5,127</sup> However, the role of NCR<sup>-</sup> ILC3s in patients with IBD remains unclear. In mice IL-17A–producing NCR<sup>-</sup> ILC3s were indispensable for the pathogenesis of mouse colitis models that resemble human

IBD, such as *Helicobacter hepaticus*–induced colitis<sup>128</sup> and microbiota-dependent colitis in *Tbx21*<sup>-/-</sup>*Rag2*<sup>-/-</sup> mice, which lack T-bet in the innate immune system.<sup>129</sup> Anti-IL-17A treatment attenuated intestinal inflammation in these models, suggesting the importance of IL-17A–producing NCR<sup>-</sup> ILC3s in the induction of inflammation in these mice. In contrast to mouse models, the frequency of NCR<sup>-</sup> ILC3s in inflamed tissues from patients with CD was comparable with that in noninflamed control tissue.<sup>30</sup> In addition, inhibition of IL-17A by the anti-IL-17A antibody secukinumab was ineffective in a clinical trial of patients with moderate-to-severe CD.<sup>130</sup> These findings suggest that NCR<sup>-</sup> ILC3s and IL-17A likely do not play a critical role in human CD.

ILC3s were also shown to have a regulatory role in the intestine through interaction with other immune cell subsets, such as macrophages and T cells. It was shown that on IL-1 $\beta$  stimulation, ILC3-derived GM-CSF drives retinoic acid and TGF- $\beta$  production from macrophages to generate Treg cells that limit intestinal inflammation (Fig 4).<sup>131</sup> MHC class II is expressed on NCR<sup>-</sup> ILC3s in both human subjects and mice. MHC class II<sup>+</sup> NCR<sup>-</sup> ILC3s suppress the response of CD4<sup>+</sup> T cells to commensal bacteria by inducing cell death. Moreover, depletion of MHC class II<sup>+</sup> ILC3s in mice results in commensal

bacteria-dependent colitis induced by increased CD4<sup>+</sup> T-cell activation.<sup>132,133</sup> These findings suggest that MHC class II<sup>+</sup> NCR<sup>−</sup> ILC3s are important for the maintenance of intestinal homeostasis by limiting bacteria-induced T-cell activation through MHC class II (Fig 4). Interestingly, the expression level of MHC class II on intestinal ILC3s from patients with pediatric CD was significantly lower than that on ILC3s in the intestines of control subjects. In addition, the expression level of MHC class II on ILC3s in the intestines of patients with pediatric CD negatively correlated with the frequency of T<sub>H</sub>17 cells.<sup>133</sup> These findings suggest that alteration of MHC class II expression on ILC3s is one of the mechanisms that controls the frequency of pathogenic T<sub>H</sub>17 cells.

Taken together, NCR<sup>−</sup> ILC3s can have both protective and pathogenic roles in patients with IBD, and further studies are needed to clarify these roles.

## CONCLUDING REMARKS

Before the discovery of ILCs, T cells were considered the main cell type responsible for the pathogenesis of inflammatory disease. However, recent studies have shown that ILCs are also essential players in many inflammatory diseases. It is now clear that ILCs play critical roles not only in the initiation of inflammation but also in chronic inflammation and resolution of inflammation through interactions with tissue cells and other immune cells. Further studies are needed to reveal the entire picture of interaction between various cell types and clarify the roles of ILCs in different types of inflammation. Findings from these studies could potentially lead to the development of new therapies for inflammatory diseases that target ILC activation pathways.

We thank Anna Głobińska (SIAF) for her assistance in drawing figures. We also thank Kafi N. Ealey for critical reading of the manuscript.

## REFERENCES

- Cherrier M, Sawa S, Eberl G. Notch, Id2, and RORgammat sequentially orchestrate the fetal development of lymphoid tissue inducer cells. *J Exp Med* 2012;209:729-40.
- Constantinides MG, McDonald BD, Verhoef PA, Bendelac A. A committed precursor to innate lymphoid cells. *Nature* 2014;508:397-401.
- Klose CS, Flach M, Mohle L, Rogell L, Hoyler T, Ebert K, et al. Differentiation of type 1 ILCs from a common progenitor to all helper-like innate lymphoid cell lineages. *Cell* 2014;157:340-56.
- Spits H, Artis D, Colonna M, Dieffenbach A, Di Santo JP, Eberl G, et al. Innate lymphoid cells—a proposal for uniform nomenclature. *Nat Rev Immunol* 2013;13:145-9.
- Bernink JH, Peters CP, Munneke M, te Velde AA, Meijer SL, Weijer K, et al. Human type 1 innate lymphoid cells accumulate in inflamed mucosal tissues. *Nat Immunol* 2013;14:221-9.
- Fuchs A, Vermi W, Lee JS, Lonardi S, Gilfillan S, Newberry RD, et al. Intraepithelial type 1 innate lymphoid cells are a unique subset of IL-12- and IL-15-responsive IFN- $\gamma$ -producing cells. *Immunity* 2013;38:769-81.
- Gordon SM, Chaix J, Rupp LJ, Wu J, Madera S, Sun JC, et al. The transcription factors T-bet and Eomes control key checkpoints of natural killer cell maturation. *Immunity* 2012;36:55-67.
- Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, et al. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature* 2010;463:540-4.
- Mjosberg JM, Trifari S, Crellin NK, Peters CP, van Drunen CM, Piet B, et al. Human IL-25- and IL-33-responsive type 2 innate lymphoid cells are defined by expression of CCR2 and CD161. *Nat Immunol* 2011;12:1055-62.
- Neill DR, Wong SH, Bellosi A, Flynn RJ, Daly M, Langford TK, et al. Nuocytes represent a new innate effector leukocyte that mediates type-2 immunity. *Nature* 2010;464:1367-70.
- Price AE, Liang HE, Sullivan BM, Reinhardt RL, Easley CJ, Erle DJ, et al. Systemically dispersed innate IL-13-expressing cells in type 2 immunity. *Proc Natl Acad Sci U S A* 2010;107:11489-94.
- Mjosberg J, Bernink J, Golebski K, Karrich JJ, Peters CP, Blom B, et al. The transcription factor GATA3 is essential for the function of human type 2 innate lymphoid cells. *Immunity* 2012;37:649-59.
- Wilhelm C, Hirota K, Stieglitz B, Van Snick J, Tolaini M, Lahl K, et al. An IL-9 fate reporter demonstrates the induction of an innate IL-9 response in lung inflammation. *Nat Immunol* 2011;12:1071-7.
- Turner JE, Morrison PJ, Wilhelm C, Wilson M, Ahlfors H, Renaud JC, et al. IL-9-mediated survival of type 2 innate lymphoid cells promotes damage control in helminth-induced lung inflammation. *J Exp Med* 2013;210:2951-65.
- Pelly VS, Kannan Y, Coomes SM, Entwistle LJ, Ruckerl D, Seddon B, et al. IL-4-producing ILC2s are required for the differentiation of T2 cells following *Heligmosomoides polygyrus* infection. *Mucosal Immunol* 2016 [Epub ahead of print].
- Yagi R, Zhong C, Northrup DL, Yu F, Bouladoux N, Spencer S, et al. The transcription factor GATA3 is critical for the development of all IL-7R $\alpha$ -expressing innate lymphoid cells. *Immunity* 2014;40:378-88.
- Serafini N, Klein Wolterink RG, Satoh-Takayama N, Xu W, Vossheerich CA, Hendriks RW, et al. Gata3 drives development of RORgammat+ group 3 innate lymphoid cells. *J Exp Med* 2014;211:199-208.
- Bal SM, Bernink JH, Nagasawa M, Groot J, Shikhaagaie MM, Golebski K, et al. IL-1 $\beta$ , IL-4 and IL-12 control the fate of group 2 innate lymphoid cells in human airway inflammation in the lungs. *Nat Immunol* 2016;17:636-45.
- Ohne Y, Silver JS, Thompson-Snipes L, Collet MA, Blanck JP, Cantarel BL, et al. IL-1 is a critical regulator of group 2 innate lymphoid cell function and plasticity. *Nat Immunol* 2016;17:646-55.
- Xue L, Salimi M, Panse I, Mjosberg JM, McKenzie AN, Spits H, et al. Prostaglandin D2 activates group 2 innate lymphoid cells through chemoattractant receptor-homologous molecule expressed on TH2 cells. *J Allergy Clin Immunol* 2014;133:1184-94.
- Doherty TA, Khorram N, Lund S, Mehta AK, Croft M, Broide DH. Lung type 2 innate lymphoid cells express cysteinyl leukotriene receptor 1, which regulates TH2 cytokine production. *J Allergy Clin Immunol* 2013;132:205-13.
- Monticelli LA, Sonnenberg GF, Abt MC, Alenghat T, Ziegler CG, Doering TA, et al. Innate lymphoid cells promote lung-tissue homeostasis after infection with influenza virus. *Nat Immunol* 2011;12:1045-54.
- Mebius RE, Rennert P, Weissman IL. Developing lymph nodes collect CD4+CD3- LT $\beta$  cells that can differentiate to APC, NK cells, and follicular cells but not T or B cells. *Immunity* 1997;7:493-504.
- Adachi S, Yoshida H, Kataoka H, Nishikawa S. Three distinctive steps in Peyer's patch formation of murine embryo. *Int Immunol* 1997;9:507-14.
- Cella M, Fuchs A, Vermi W, Facchetti F, Otero K, Lennerz JK, et al. A human natural killer cell subset provides an innate source of IL-22 for mucosal immunity. *Nature* 2009;457:722-5.
- Satoh-Takayama N, Lesjean-Pottier S, Vieira P, Sawa S, Eberl G, Vossheerich CA, et al. IL-7 and IL-15 independently program the differentiation of intestinal CD3-NKp46+ cell subsets from Id2-dependent precursors. *J Exp Med* 2010;207:273-80.
- Sonnenberg GF, Monticelli LA, Elloso MM, Fouser LA, Artis D. CD4(+) lymphoid tissue-inducer cells promote innate immunity in the gut. *Immunity* 2011;34:122-34.
- Vonarbourg C, Mortha A, Bui VL, Hernandez PP, Kiss EA, Hoyler T, et al. Regulated expression of nuclear receptor RORgammat confers distinct functional fates to NK cell receptor-expressing RORgammat(+) innate lymphocytes. *Immunity* 2010;33:736-51.
- Crellin NK, Trifari S, Kaplan CD, Satoh-Takayama N, Di Santo JP, Spits H. Regulation of cytokine secretion in human CD127(+) LT $\alpha$ -like innate lymphoid cells by Toll-like receptor 2. *Immunity* 2010;33:752-64.
- Bernink JH, Krabbendam L, Germar K, de Jong E, Gronke K, Kofoed-Nielsen M, et al. Interleukin-12 and -23 control plasticity of CD127(+) group 1 and group 3 innate lymphoid cells in the intestinal lamina propria. *Immunity* 2015;43:146-60.
- Mizuno S, Mikami Y, Kamada N, Handa T, Hayashi A, Sato T, et al. Cross-talk between RORgammat+ innate lymphoid cells and intestinal macrophages induces mucosal IL-22 production in Crohn's disease. *Inflamm Bowel Dis* 2014;20:1426-34.
- Lim AI, Menegatti S, Bustamante J, Le Bourhis L, Allez M, Rogge L, et al. IL-12 drives functional plasticity of human group 2 innate lymphoid cells. *J Exp Med* 2016;213:569-83.
- Silver JS, Kearley J, Copenhaver AM, Sanden C, Mori M, Yu L, et al. Inflammatory triggers associated with exacerbations of COPD orchestrate plasticity of group 2 innate lymphoid cells in the lungs. *Nat Immunol* 2016;17:626-35.
- Grotenboer NS, Ketelaar ME, Koppelman GH, Nawijn MC. Decoding asthma: translating genetic variation in IL33 and IL1RL1 into disease pathophysiology. *J Allergy Clin Immunol* 2013;131:856-65.

35. Tamari M, Tanaka S, Hirota T. Genome-wide association studies of allergic diseases. *Allergol Int* 2013;62:21-8.
36. Oboki K, Ohno T, Kajiwara N, Saito H, Nakae S. IL-33 and IL-33 receptors in host defense and diseases. *Allergol Int* 2010;59:143-60.
37. Bartemes KR, Kephart GM, Fox SJ, Kita H. Enhanced innate type 2 immune response in peripheral blood from patients with asthma. *J Allergy Clin Immunol* 2014;134:671-8.e4.
38. Liu T, Wu J, Zhao J, Wang J, Zhang Y, Liu L, et al. Type 2 innate lymphoid cells: a novel biomarker of eosinophilic airway inflammation in patients with mild to moderate asthma. *Respir Med* 2015;109:1391-6.
39. Smith SG, Chen R, Kjarsgaard M, Huang C, Oliveria JP, O'Byrne PM, et al. Increased numbers of activated group 2 innate lymphoid cells in the airways of patients with severe asthma and persistent airway eosinophilia. *J Allergy Clin Immunol* 2016;137:75-86.e8.
40. Christianson CA, Goplen NP, Zafar I, Irvin C, Good JT Jr, Rollins DR, et al. Persistence of asthma requires multiple feedback circuits involving type 2 innate lymphoid cells and IL-33. *J Allergy Clin Immunol* 2015;136:59-68.e14.
41. Nagakumar P, Denney L, Fleming L, Bush A, Lloyd CM, Saglani S. Type 2 innate lymphoid cells in induced sputum from children with severe asthma. *J Allergy Clin Immunol* 2016;137:624-6.e6.
42. Morita H, Arae K, Unno H, Miyauchi K, Toyama S, Nambu A, et al. An interleukin-33-mast cell-interleukin-2 axis suppresses papain-induced allergic inflammation by promoting regulatory T cell numbers. *Immunity* 2015;43:175-86.
43. Kabata H, Moro K, Fukunaga K, Suzuki Y, Miyata J, Masaki K, et al. Thymic stromal lymphopoietin induces corticosteroid resistance in natural helper cells during airway inflammation. *Nat Commun* 2013;4:2675.
44. Halim TY, Krauss RH, Sun AC, Takei F. Lung natural helper cells are a critical source of Th2 cell-type cytokines in protease allergen-induced airway inflammation. *Immunity* 2012;36:451-63.
45. Barlow JL, Bellosi A, Hardman CS, Drynan LF, Wong SH, Cruickshank JP, et al. Innate IL-13-producing nuocytes arise during allergic lung inflammation and contribute to airways hyperactivity. *J Allergy Clin Immunol* 2012;129:191-8, e1-4.
46. Klein Wolterink RG, Kleijne A, van Nimwegen M, Bergen I, de Bruijn M, Levani Y, et al. Pulmonary innate lymphoid cells are major producers of IL-5 and IL-13 in murine models of allergic asthma. *Eur J Immunol* 2012;42:1106-16.
47. Bartemes KR, Iijima K, Kobayashi T, Kephart GM, McKenzie AN, Kita H. IL-33-responsive lineage- CD25+ CD44(hi) lymphoid cells mediate innate type 2 immunity and allergic inflammation in the lungs. *J Immunol* 2012;188:1503-13.
48. Doherty TA, Khorram N, Chang JE, Kim HK, Rosenthal P, Croft M, et al. STAT6 regulates natural helper cell proliferation during lung inflammation initiated by *Alternaria*. *Am J Physiol Lung Cell Mol Physiol* 2012;303:L577-88.
49. Chang YJ, Kim HY, Albacker LA, Baumgarth N, McKenzie AN, Smith DE, et al. Innate lymphoid cells mediate influenza-induced airway hyper-reactivity independently of adaptive immunity. *Nat Immunol* 2011;12:631-8.
50. Takai T, Ikeda S. Barrier dysfunction caused by environmental proteases in the pathogenesis of allergic diseases. *Allergol Int* 2011;60:25-35.
51. Milne J, Brand S. Occupational asthma after inhalation of dust of the proteolytic enzyme, papain. *Br J Ind Med* 1975;32:302-7.
52. Oboki K, Ohno T, Kajiwara N, Arae K, Morita H, Ishii A, et al. IL-33 is a crucial amplifier of innate rather than acquired immunity. *Proc Natl Acad Sci U S A* 2010;107:18581-6.
53. Motomura Y, Morita H, Moro K, Nakae S, Artis D, Endo TA, et al. Basophil-derived interleukin-4 controls the function of natural helper cells, a member of ILC2s, in lung inflammation. *Immunity* 2014;40:758-71.
54. Maazi H, Patel N, Sankaranarayanan I, Suzuki Y, Rigas D, Soroosh P, et al. ICOS: ICOS-ligand interaction is required for type 2 innate lymphoid cell function, homeostasis, and induction of airway hyperactivity. *Immunity* 2015;42:538-51.
55. Morita H, Saito H, Matsumoto K, Nakae S. Regulatory roles of mast cells in immune responses. *Semin Immunopathol* 2016;38:623-9.
56. Shaw JL, Fakhri S, Citardi MJ, Porter PC, Corry DB, Kheradmand F, et al. IL-33-responsive innate lymphoid cells are an important source of IL-13 in chronic rhinosinusitis with nasal polyps. *Am J Respir Crit Care Med* 2013;188:432-9.
57. Reese TA, Liang HE, Tager AM, Luster AD, Van Rooijen N, Voehringer D, et al. Chitin induces accumulation in tissue of innate immune cells associated with allergy. *Nature* 2007;447:92-6.
58. Van Dyken SJ, Mohapatra A, Nussbaum JC, Molofsky AB, Thornton EE, Ziegler SF, et al. Chitin activates parallel immune modules that direct distinct inflammatory responses via innate lymphoid type 2 and gammadelta T cells. *Immunity* 2014;40:414-24.
59. Stier MT, Bloodworth MH, Toki S, Newcomb DC, Goleniewska K, Boyd KL, et al. Respiratory syncytial virus infection activates IL-13-producing group 2 innate lymphoid cells through thymic stromal lymphopoietin. *J Allergy Clin Immunol* 2016;138:814-24.e11.
60. Lee HC, Headley MB, Loo YM, Berlin A, Gale M Jr, Debley JS, et al. Thymic stromal lymphopoietin is induced by respiratory syncytial virus-infected airway epithelial cells and promotes a type 2 response to infection. *J Allergy Clin Immunol* 2012;130:1187-96.e5.
61. Hong JY, Bentley JK, Chung Y, Lei J, Steenrod JM, Chen Q, et al. Neonatal rhinovirus induces mucous metaplasia and airways hyperresponsiveness through IL-25 and type 2 innate lymphoid cells. *J Allergy Clin Immunol* 2014;134:429-39.
62. Jackson DJ, Gangnon RE, Evans MD, Roberg KA, Anderson EL, Pappas TE, et al. Wheezing rhinovirus illnesses in early life predict asthma development in high-risk children. *Am J Respir Crit Care Med* 2008;178:667-72.
63. Gold MJ, Antignano F, Halim TY, Hirota JA, Blanchet MR, Zaph C, et al. Group 2 innate lymphoid cells facilitate sensitization to local, but not systemic, TH2-inducing allergen exposures. *J Allergy Clin Immunol* 2014;133:1142-8.
64. Halim TY, Steer CA, Matha L, Gold MJ, Martinez-Gonzalez I, McNagny KM, et al. Group 2 innate lymphoid cells are critical for the initiation of adaptive T helper 2 cell-mediated allergic lung inflammation. *Immunity* 2014;40:425-35.
65. Drake LY, Iijima K, Kita H. Group 2 innate lymphoid cells and CD4+ T cells cooperate to mediate type 2 immune response in mice. *Allergy* 2014;69:1300-7.
66. Mirchandani AS, Besnard AG, Yip E, Scott C, Bain CC, Cerovic V, et al. Type 2 innate lymphoid cells drive CD4+ Th2 cell responses. *J Immunol* 2014;192:2442-8.
67. Halim TY, Hwang YY, Scanlon ST, Zaghouani H, Garbi N, Fallon PG, et al. Group 2 innate lymphoid cells license dendritic cells to potentiate memory TH2 cell responses. *Nat Immunol* 2016;17:57-64.
68. Barnig C, Cernadas M, Dutile S, Liu X, Perrella MA, Kazani S, et al. Lipoxin A4 regulates natural killer cell and type 2 innate lymphoid cell activation in asthma. *Sci Transl Med* 2013;5:174ra26.
69. Chang JE, Doherty TA, Baum R, Broide D. Prostaglandin D2 regulates human type 2 innate lymphoid cell chemotaxis. *J Allergy Clin Immunol* 2014;133:899-901.e3.
70. Schleimer RP, Fox CC, Naclerio RM, Plaut M, Creticos PS, Togias AG, et al. Role of human basophils and mast cells in the pathogenesis of allergic diseases. *J Allergy Clin Immunol* 1985;76:369-74.
71. Harvina IT, Levi-Schaffer F, Draber P, Friedman S, Polakovicova I, Gibbs BF, et al. Molecular targets on mast cells and basophils for novel therapies. *J Allergy Clin Immunol* 2014;134:530-44.
72. Allakhverdi Z, Smith DE, Comeau MR, Delespesse G. Cutting edge: the ST2 ligand IL-33 potently activates and drives maturation of human mast cells. *J Immunol* 2007;179:2051-4.
73. Ho LH, Ohno T, Oboki K, Kajiwara N, Suto H, Iikura M, et al. IL-33 induces IL-13 production by mouse mast cells independently of IgE-FcεpsilonRI signals. *J Leukoc Biol* 2007;82:1481-90.
74. Malo JL, Ghezzi H, D'Aquino C, L'Archeveque J, Cartier A, Chan-Yeung M. Natural history of occupational asthma: relevance of type of agent and other factors in the rate of development of symptoms in affected subjects. *J Allergy Clin Immunol* 1992;90:937-44.
75. Moro K, Kabata H, Tanabe M, Koga S, Takeno N, Mochizuki M, et al. Interferon and IL-27 antagonize the function of group 2 innate lymphoid cells and type 2 innate immune responses. *Nat Immunol* 2016;17:76-86.
76. Duerr CU, McCarthy CD, Mindt BC, Rubio M, Meli AP, Pothlichet J, et al. Type I interferon restricts type 2 immunopathology through the regulation of group 2 innate lymphoid cells. *Nat Immunol* 2016;17:65-75.
77. Molofsky AB, Van Gool F, Liang HE, Van Dyken SJ, Nussbaum JC, Lee J, et al. Interleukin-33 and interferon-gamma counter-regulate group 2 innate lymphoid cell activation during immune perturbation. *Immunity* 2015;43:161-74.
78. McHedlidze T, Kindermann M, Neves AT, Voehringer D, Neurath MF, Wirtz S. IL-27 suppresses type 2 immune responses in vivo via direct effects on group 2 innate lymphoid cells. *Mucosal Immunol* 2016 [Epub ahead of print].
79. Zhou W, Toki S, Zhang J, Goleniewska K, Newcomb DC, Cephus JY, et al. Prostaglandin I2 signaling and inhibition of group 2 innate lymphoid cell responses. *Am J Respir Crit Care Med* 2016;193:31-42.
80. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med* 2012;18:716-25.
81. Julia V, Macia L, Dombrowicz D. The impact of diet on asthma and allergic diseases. *Nat Rev Immunol* 2015;15:308-22.
82. Kim HY, Lee HJ, Chang YJ, Pichavant M, Shore SA, Fitzgerald KA, et al. Interleukin-17-producing innate lymphoid cells and the NLRP3 inflammasome facilitate obesity-associated airway hyperactivity. *Nat Med* 2014;20:54-61.
83. Everaere L, Ait-Yahia S, Molendi-Coste O, Vorm H, Quemener S, LeVu P, et al. Innate lymphoid cells contribute to allergic airway disease exacerbation by obesity. *J Allergy Clin Immunol* 2016 [Epub ahead of print].
84. Brestoff JR, Kim BS, Saenz SA, Stine RR, Monticelli LA, Sonnenberg GF, et al. Group 2 innate lymphoid cells promote beiging of white adipose tissue and limit obesity. *Nature* 2015;519:242-6.
85. Weidinger S, Novak N. Atopic dermatitis. *Lancet* 2016;387:1109-22.



86. Soumelis V, Reche PA, Kanzler H, Yuan W, Edward G, Homey B, et al. Human epithelial cells trigger dendritic cell mediated allergic inflammation by producing TSLP. *Nat Immunol* 2002;3:673-80.
87. Wang YH, Angkasekwinai P, Lu N, Voo KS, Arima K, Hanabuchi S, et al. IL-25 augments type 2 immune responses by enhancing the expansion and functions of TSLP-DC-activated Th2 memory cells. *J Exp Med* 2007;204:1837-47.
88. Savinko T, Matikainen S, Saarialho-Kere U, Lehto M, Wang G, Lehtimäki S, et al. IL-33 and ST2 in atopic dermatitis: expression profiles and modulation by triggering factors. *J Invest Dermatol* 2012;132:1392-400.
89. Kim BS, Siracusa MC, Saenz SA, Noti M, Monticelli LA, Sonnenberg GF, et al. TSLP elicits IL-33-independent innate lymphoid cell responses to promote skin inflammation. *Sci Transl Med* 2013;5:170ra16.
90. Salimi M, Barlow JL, Saunders SP, Xue L, Gutowska-Owsiak D, Wang X, et al. A role for IL-25 and IL-33-driven type-2 innate lymphoid cells in atopic dermatitis. *J Exp Med* 2013;210:2939-50.
91. Teunissen MB, Munneke JM, Bernink JH, Spuls PI, Res PC, Te Velde A, et al. Composition of innate lymphoid cell subsets in the human skin: enrichment of NCR(+) ILC3 in lesional skin and blood of psoriasis patients. *J Invest Dermatol* 2014;134:2351-60.
92. Imai Y, Yasuda K, Sakaguchi Y, Haneda T, Mizutani H, Yoshimoto T, et al. Skin-specific expression of IL-33 activates group 2 innate lymphoid cells and elicits atopic dermatitis-like inflammation in mice. *Proc Natl Acad Sci U S A* 2013;110:13921-6.
93. Li M, Messaddeq N, Teletin M, Pasquali JL, Metzger D, Chambon P. Retinoid X receptor ablation in adult mouse keratinocytes generates an atopic dermatitis triggered by thymic stromal lymphopoietin. *Proc Natl Acad Sci U S A* 2005;102:14795-800.
94. Yoo J, Omori M, Gyarmati D, Zhou B, Aye T, Brewer A, et al. Spontaneous atopic dermatitis in mice expressing an inducible thymic stromal lymphopoietin transgene specifically in the skin. *J Exp Med* 2005;202:541-9.
95. Kim BS, Wang K, Siracusa MC, Saenz SA, Brestoff JR, Monticelli LA, et al. Basophils promote innate lymphoid cell responses in inflamed skin. *J Immunol* 2014;193:3717-25.
96. Otsuka A, Nonomura Y, Kabashima K. Roles of basophils and mast cells in cutaneous inflammation. *Semin Immunopathol* 2016;38:563-70.
97. Hoyler T, Klose CS, Souabni A, Turqueti-Neves A, Pfeifer D, Rawlins EL, et al. The transcription factor GATA-3 controls cell fate and maintenance of type 2 innate lymphoid cells. *Immunity* 2012;37:634-48.
98. Tang A, Amagai M, Granger LG, Stanley JR, Udey MC. Adhesion of epidermal Langerhans cells to keratinocytes mediated by E-cadherin. *Nature* 1993;361:82-5.
99. Trautmann A, Altnauer F, Akdis M, Simon HU, Disch R, Brocker EB, et al. The differential fate of cadherins during T-cell-induced keratinocyte apoptosis leads to spongiolysis in eczematous dermatitis. *J Invest Dermatol* 2001;117:927-34.
100. Nestle FO, Kaplan DH, Barker J. Psoriasis. *N Engl J Med* 2009;361:496-509.
101. Bowcock AM, Krueger JG. Getting under the skin: the immunogenetics of psoriasis. *Nat Rev Immunol* 2005;5:699-711.
102. Villanova F, Flutter B, Tosi I, Grys K, Sreeneebus H, Perera GK, et al. Characterization of innate lymphoid cells in human skin and blood demonstrates increase of NKp44+ ILC3 in psoriasis. *J Invest Dermatol* 2014;134:984-91.
103. Dyring-Andersen B, Geisler C, Agerbeck C, Lauritsen JP, Gudjonsdottir SD, Skov L, et al. Increased number and frequency of group 3 innate lymphoid cells in non-lesional psoriatic skin. *Br J Dermatol* 2014;170:609-16.
104. Pantelyushin S, Haak S, Ingold B, Kulig P, Heppner FL, Navarini AA, et al. RORgammat+ innate lymphocytes and gammadelta T cells initiate psoriasisiform plaque formation in mice. *J Clin Invest* 2012;122:2252-6.
105. Li Z, Hodgkinson T, Gothard EJ, Boroumand S, Lamb R, Cummins I, et al. Epidermal Notch1 recruits RORgamma(+) group 3 innate lymphoid cells to orchestrate normal skin repair. *Nat Commun* 2016;7:11394.
106. Rothenberg ME. Eosinophilic gastrointestinal disorders (EGID). *J Allergy Clin Immunol* 2004;113:11-29.
107. Cianferoni A, Spigelman JM. Eosinophilic esophagitis and gastroenteritis. *Curr Allergy Asthma Rep* 2015;15:58.
108. Mishra A, Hogan SP, Brandt EB, Rothenberg ME. IL-5 promotes eosinophil trafficking to the esophagus. *J Immunol* 2002;168:2464-9.
109. Blanchard C, Mingler MK, Vicario M, Abonia JP, Wu YY, Lu TX, et al. IL-13 involvement in eosinophilic esophagitis: transcriptome analysis and reversibility with glucocorticoids. *J Allergy Clin Immunol* 2007;120:1292-300.
110. Rothenberg ME, Spigelman JM, Sherrill JD, Annaiah K, Martin LJ, Cianferoni A, et al. Common variants at 5q22 associate with pediatric eosinophilic esophagitis. *Nat Genet* 2010;42:289-91.
111. Simon D, Radonjic-Hosli S, Straumann A, Yousefi S, Simon HU. Active eosinophilic esophagitis is characterized by epithelial barrier defects and eosinophil extracellular trap formation. *Allergy* 2015;70:443-52.
112. Doherty TA, Baum R, Newbury RO, Yang T, Dohil R, Aquino M, et al. Group 2 innate lymphocytes (ILC2) are enriched in active eosinophilic esophagitis. *J Allergy Clin Immunol* 2015;136:792-4.e3.
113. Shoda T, Matsuda A, Arai K, Shimizu H, Morita H, Orihara K, et al. Sera of patients with infantile eosinophilic gastroenteritis showed a specific increase in both thymic stromal lymphopoietin and IL-33 levels. *J Allergy Clin Immunol* 2016;138:299-303.
114. Neurath MF. Cytokines in inflammatory bowel disease. *Nat Rev Immunol* 2014;14:329-42.
115. Jostins L, Ripke S, Weersma RK, Duerr RH, McGovern DP, Hui KY, et al. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012;491:119-24.
116. Fuss IJ, Neurath M, Boirivant M, Klein JS, de la Motte C, Strong SA, et al. Disparate CD4+ lamina propria (LP) lymphokine secretion profiles in inflammatory bowel disease. Crohn's disease LP cells manifest increased secretion of IFN-gamma, whereas ulcerative colitis LP cells manifest increased secretion of IL-5. *J Immunol* 1996;157:1261-70.
117. Sakuraba A, Sato T, Kamada N, Kitazume M, Sugita A, Hibi T. Th1/Th17 immune response is induced by mesenteric lymph node dendritic cells in Crohn's disease. *Gastroenterology* 2009;137:1736-45.
118. Rovedatti L, Kudo T, Biancheri P, Sarra M, Knowles CH, Rampton DS, et al. Differential regulation of interleukin 17 and interferon gamma production in inflammatory bowel disease. *Gut* 2009;58:1629-36.
119. Takayama T, Kamada N, Chinen H, Okamoto S, Kitazume MT, Chang J, et al. Imbalance of NKp44(+)NKp46(-) and NKp44(-)NKp46(+) natural killer cells in the intestinal mucosa of patients with Crohn's disease. *Gastroenterology* 2010;139:882-92, e1-3.
120. Cella M, Otero K, Colonna M. Expansion of human NK-22 cells with IL-7, IL-2, and IL-1beta reveals intrinsic functional plasticity. *Proc Natl Acad Sci U S A* 2010;107:10961-6.
121. Pickert G, Neufert C, Leppkes M, Zheng Y, Wittkopf N, Warntjen M, et al. STAT3 links IL-22 signaling in intestinal epithelial cells to mucosal wound healing. *J Exp Med* 2009;206:1465-72.
122. Kim CJ, Nazli A, Rojas OL, Chege D, Alidina Z, Huibner S, et al. A role for mucosal IL-22 production and Th22 cells in HIV-associated mucosal immunopathogenesis. *Mucosal Immunol* 2012;5:670-80.
123. Mizoguchi A. Healing of intestinal inflammation by IL-22. *Inflamm Bowel Dis* 2012;18:1777-84.
124. Bishop JL, Roberts ME, Beer JL, Huang M, Chehal MK, Fan X, et al. Lyn activity protects mice from DSS colitis and regulates the production of IL-22 from innate lymphoid cells. *Mucosal Immunol* 2014;7:405-16.
125. Sawa S, Lochner M, Satoh-Takayama N, Dulauroy S, Berard M, Kleinschek M, et al. RORgammat+ innate lymphoid cells regulate intestinal homeostasis by integrating negative signals from the symbiotic microbiota. *Nat Immunol* 2011;12:320-6.
126. Eken A, Singh AK, Treuting PM, Oukka M. IL-23R+ innate lymphoid cells induce colitis via interleukin-22-dependent mechanism. *Mucosal Immunol* 2014;7:143-54.
127. Geremia A, Arancibia-Carcamo CV, Fleming MP, Rust N, Singh B, Mortensen NJ, et al. IL-23-responsive innate lymphoid cells are increased in inflammatory bowel disease. *J Exp Med* 2011;208:1127-33.
128. Buonocore S, Ahern PP, Uhlig HH, Ivanov II, Littman DR, Maloy KJ, et al. Innate lymphoid cells drive interleukin-23-dependent intestinal pathology. *Nature* 2010;464:1371-5.
129. Powell N, Walker AW, Stolarczyk E, Canavan JB, Gokmen MR, Marks E, et al. The transcription factor T-bet regulates intestinal inflammation mediated by interleukin-7 receptor+ innate lymphoid cells. *Immunity* 2012;37:674-84.
130. Hueber W, Sands BE, Lewitzky S, Vandemeulebroeck M, Reinisch W, Higgins PD, et al. Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate to severe Crohn's disease: unexpected results of a randomised, double-blind placebo-controlled trial. *Gut* 2012;61:1693-700.
131. Mortha A, Chudnovskiy A, Hashimoto D, Bogunovic M, Spencer SP, Belkaid Y, et al. Microbiota-dependent crosstalk between macrophages and ILC3 promotes intestinal homeostasis. *Science* 2014;343:1249288.
132. Hepworth MR, Monticelli LA, Fung TC, Ziegler CG, Grunberg S, Sinha R, et al. Innate lymphoid cells regulate CD4+ T-cell responses to intestinal commensal bacteria. *Nature* 2013;498:113-7.
133. Hepworth MR, Fung TC, Masur SH, Kelsen JR, McConnell FM, Dubrot J, et al. Immune tolerance. Group 3 innate lymphoid cells mediate intestinal selection of commensal bacteria-specific CD4(+) T cells. *Science* 2015;348:1031-5.